

FACIES ARCHITECTURE, GEOCHEMISTRY AND TECTONIC
SIGNIFICANCE OF THE URAL VOLCANICS AND THE MOUNT
HOPE VOLCANICS, CENTRAL LACHLAN OROGEN, NSW

Katharine F. Bull
M.Sc. University of Alaska Fairbanks



Submitted in fulfilment of the requirements
for the degree of Doctor of Philosophy


University of Tasmania
Australia
May, 2006

STATEMENT AND AUTHORITY OF ACCESS

This thesis contains no material which has been accepted for a degree or diploma by the University or any other institution and, the best of my knowledge and belief, no material previously published or written by another person except where due acknowledgement is made in the text of this thesis.

This thesis may be made available for loan and limited copying in accordance with the *Copyright Act* of 1968.

Date: 12 May, 2006

A handwritten signature in blue ink, appearing to read 'Katharine F. Bull', with a long vertical stroke extending downwards from the end of the signature.

Katharine F. Bull

ABSTRACT

This study focuses on facies analysis, geochemistry, geochronology and tectonic significance of the Ural Volcanics (UV) and Mount Hope Volcanics (MHV) in the Central Lachlan Orogen in New South Wales. The UV and MHV overlie non-volcanic sedimentary, below wave base, submarine facies within two intracontinental rift basins, the Rast and Mount Hope Troughs. The UV and MHV consist primarily of felsic, coherent facies and associated felsic monomictic breccia facies. These volcanic facies are interpreted to represent submarine lava-sill complexes, which define intrabasinal, effusive, volcanic and shallow intrusive centres. The UV include at least 35 separate lava or sill emplacement units that amount to $\sim 10 \text{ km}^3$. In the MHV, at least 18 lavas and sills are present, and have thicknesses up to $\sim 120 \text{ m}$. The combined volume of the two largest MHV units is estimated to be $\leq 1.5 \text{ km}^3$.

In the UV, siltstone-matrix monomictic breccia facies is characterised by continuously laminated siltstone matrix between monomictic, non-vesicular, felsic clasts. This facies is interpreted to form from water-settled sediment deposited between lava clasts, and must therefore occur on the upper margin of a lava. The presence of conformable, continuous laminae helps to distinguish this facies from peperite. Hence, correct identification of the siltstone-matrix monomictic breccia facies is critical in distinguishing lavas from sills.

The autoclastic facies in the UV and MHV account for up to 10% of single emplacement units. In most cases, the clasts have blocky or slabby shapes and are flow-banded, implying that autobrecciation was the main fragmentation mechanism. Neither *in situ* or resedimented hyaloclastite are recognised in the UV or MHV, in contrast to other submarine felsic lavas and domes elsewhere.

Pumice-rich volcanoclastic facies in the UV and MHV are less voluminous than the coherent and monomictic breccia facies. The UV pumice-rich facies are interpreted to represent felsic pyroclastic facies erupted in a single, open-vent explosive eruption from a local vent and transported in, and deposited from, submarine gravity currents (pumice-rich facies association),

and settled from suspension in the water column (fiamme-siltstone breccia facies). Similar syn-eruptive pyroclastic facies occur in the MHV, but their source has not been identified.

Fiamme-bearing facies occur in both the UV and MHV, in fiamme-bearing pyroclastic facies and as pseudoclastic textures. Fiamme textures can also be formed in a variety of other ways. The common genetic use of the term fiamme for textures produced by welding compaction is easily misinterpreted. 'Fiamme' would be better used descriptively to mean elongate lenses or domains of the same mineralogy, texture or composition, which define a pre-tectonic foliation, and are separated by domains of different mineralogy, texture or composition.

Results from LA-ICPMS U/Pb dating of zircons indicate that the UV and the MHV were roughly coeval and erupted in the Late Silurian-Early Devonian, within the time period ~420-410 Ma. Both successions consist of dacites and rhyolites and have A-type to transitional I-type geochemical affinities. High-level, felsic, A-type plutons with myrmekitic and/or granophyric texture occur in both study areas. Geochemistry suggests they are comagmatic with the volcanic facies. Cross-cutting mafic to intermediate dykes and small intrusions are not comagmatic with the felsic coherent facies.

A modern analogue for the UV and MHV felsic has not been recognised. The closest analogue is the Late Devonian-Early Carboniferous Iberian Pyrite Belt (IPB). Parts of the IPB contain similar felsic coherent, monomictic breccia facies, and syn-eruptive pyroclastic facies as the UV and MHV. The IPB volcanic rocks are also A-type in composition. The numerous similarities of the IPB to the UV and MHV suggest the Australian successions have great potential for hosting volcanic-hosted massive sulfide (VHMS) deposits, however, neither previous exploration nor mapping during this study have uncovered any VHMS-related altered zones or prospects.

TABLE OF CONTENTS

STATEMENT AND AUTHORITY OF ACCESS	ii
ABSTRACT	iii
ACKNOWLEDGEMENTS	xi
CHAPTER 1. INTRODUCTION	
1.1 INTRODUCTION	1
1.2 SIGNIFICANCE AND AIMS	2
1.3 LOCATION AND ACCESS	5
1.4 METHODS AND APPROACH	7
1.5 THESIS ORGANISATION	8
CHAPTER 2. REGIONAL SETTING OF THE URAL VOLCANICS AND THE MOUNT HOPE VOLCANICS IN THE LACHLAN OROGEN	
2.1 INTRODUCTION	10
2.2 LACHLAN OROGEN	10
2.3 ORDOVICIAN TO CARBONIFEROUS EVOLUTION OF THE LACHLAN OROGEN	13
<i>Ordovician-Silurian</i>	13
<i>Silurian-Devonian</i>	16
<i>Silurian-Devonian Magmatism</i>	18
<i>Middle Devonian-Carboniferous</i>	22
<i>Mineral Deposits Associated with the Lachlan Orogen</i>	26
2.4 SILURO-DEVONIAN VOLCANIC SUCCESSIONS IN THE LACHLAN OROGEN	27
2.5 REGIONAL GEOLOGY OF THE URAL RANGE AND MOUNT HOPE AREAS	28
<i>Ural Range Study Area</i>	28
<i>Mount Hope Study Area</i>	37

CHAPTER 3. FACIES AND FACIES ASSOCIATIONS OF THE URAL VOLCANICS, URAL RANGE AREA

3.1 INTRODUCTION	43
3.2 REGIONAL GEOLOGY OF THE UV	43
<i>Structure</i>	46
<i>Diagenesis and Regional Metamorphism</i>	48
<i>Mineralisation</i>	52
3.3 FACIES AND FACIES ASSOCIATIONS OF THE URAL VOLCANICS	53
<i>Felsic Coherent Facies Association</i>	58
<u><i>Felsic Coherent Facies</i></u>	58
<u><i>Monomictic Breccia Facies</i></u>	65
<u><i>Siltstone-matrix Monomictic Breccia Facies</i></u>	66
<u><i>Interpretation</i></u>	68
<u><i>Tabular Felsic Coherent Facies</i></u>	72
<u><i>Interpretation</i></u>	72
<u><i>Dacite Porphyry Facies</i></u>	73
<u><i>Interpretation</i></u>	74
<i>Mafic-Intermediate Coherent Facies Association</i>	74
<u><i>Basalt to Andesite Facies</i></u>	74
<u><i>Diorite to Leucogabbro Facies</i></u>	74
<u><i>Interpretation</i></u>	75
<i>Rhyolite–dacite–siltstone Breccia Facies</i>	75
<u><i>Interpretation</i></u>	76
<i>Fiamme–siltstone Breccia Facies</i>	77
<u><i>Interpretation</i></u>	78
<i>Pumice-rich Facies Association</i>	80
<u><i>Coarse Speckled Breccia Facies</i></u>	80
<u><i>Speckled Breccia Facies</i></u>	81
<u><i>Pumice–crystal–shard Sandstone Facies</i></u>	82
<u><i>Interpretation</i></u>	82
<i>Mixed Volcanic–Sedimentary Facies</i>	84
<u><i>Polymictic Conglomerate Facies</i></u>	84
<u><i>Interpretation</i></u>	86
<i>Non-volcanic Sedimentary Facies</i>	86
<u><i>Black Mudstone Facies</i></u>	87
<u><i>Micaceous Quartz Sandstone and Foliated Mudstone Facies</i></u>	87
<u><i>Interpretation</i></u>	88
<i>Strongly Foliated Facies</i>	89
<u><i>Interpretation</i></u>	89

CHAPTER 4. FACIES ARCHITECTURE OF THE URAL VOLCANICS	91
4.1 INTRODUCTION	91
4.2 ENVIRONMENT OF DEPOSITION OF THE URAL VOLCANICS	91
4.3 FACIES DISTRIBUTION AND ARCHITECTURE OF THE URAL VOLCANICS	95
<i>Lavas and Sills</i>	95
<i>Non-autoclastic Volcaniclastic Facies</i>	103
<i>Non-volcanic Sedimentary Facies</i>	105
4.4 STYLES OF VOLCANISM AND LOCATION OF VOLCANIC CENTRES IN THE URAL VOLCANICS	105
4.5 EVOLUTION OF THE UV IN THE URAL RANGE	109
4.6 ANALOGOUS FELSIC SUBMARINE SUCCESSIONS	112
4.7 CONCLUSIONS	113
 CHAPTER 5 - VOLCANIC, SEDIMENTARY AND PLUTONIC FACIES AND FACIES ASSOCIATIONS IN THE MOUNT HOPE AREA	
5.1 INTRODUCTION	115
5.2 REGIONAL GEOLOGY	115
<i>Structure and Metamorphism</i>	116
<i>Alteration and Mineralisation</i>	120
5.3 FACIES AND FACIES ASSOCIATIONS IN THE MOUNT HOPE AREA	121
<i>Felsic Coherent Facies Association</i>	127
<u><i>Felsic Coherent Facies</i></u>	127
<u><i>Monomictic Breccia Facies</i></u>	133
<u><i>Clast-bearing Rhyolite Facies</i></u>	136
<u><i>Interpretation</i></u>	136
<u><i>Intergrowth textures in MHV felsic coherent facies</i></u>	140
<i>Fiamme Breccia Facies Association</i>	141
<u><i>Coarse Fiamme Breccia Facies</i></u>	141
<u><i>Fine Fiamme Breccia Facies</i></u>	142
<u><i>Crystal-Shard Mudstone Facies</i></u>	145
<u><i>Interpretation</i></u>	145
<i>Rhyolite-Pumice Breccia Facies Association</i>	147
<i>Coarse Rhyolite-Pumice Breccia Facies</i>	147
<u><i>Rhyolite-Lithic-Pumice Breccia Facies</i></u>	147

<i><u>Fine Rhyolite–Pumice Breccia Facies</u></i>	151
<i><u>Crystal-Pumice Sandstone Facies</u></i>	151
<i><u>Interpretation</u></i>	153
Polymictic Volcanic-Sedimentary Conglomerate Facies	154
<i><u>Interpretation</u></i>	154
Non-Volcanic Sedimentary Facies Association	156
<i><u>Mudstone</u></i>	156
<i><u>Micaceous Quartz Sandstone</u></i>	159
<i><u>Non-volcanic Polymictic Conglomerate</u></i>	159
<i><u>Interpretation</u></i>	160
Mafic-intermediate Coherent Facies Association	161
<i><u>Basalt-Andesite Facies</u></i>	161
<i><u>Interpretation</u></i>	161
Plutonic Facies	161
<i><u>Interpretation</u></i>	163
5.4 Summary	165
 CHAPTER 6 - FACIES ARCHITECTURE OF THE MOUNT HOPE VOLCANICS	
6.1 INTRODUCTION	166
6.2 ENVIRONMENT OF DEPOSITION OF THE MOUNT HOPE VOLCANICS	166
6.3 FACIES DISTRIBUTION AND ARCHITECTURE OF THE MHV	168
6.4 STYLES OF VOLCANISM AND LOCATION OF VOLCANIC CENTRES IN THE MOUNT HOPE VOLCANICS	173
6.5 EVOLUTION OF THE MOUNT HOPE VOLCANICS	174
6.5 SUMMARY	178
 CHAPTER 7 - FIAMME TEXTURES IN VOLCANIC SUCCESSIONS: FLAMING ISSUES WITH DEFINITION AND INTERPRETATION	
7.1 INTRODUCTION	179
7.2 PREVIOUS USE OF 'FIAMME'	180
7.3 FIAMME IN WELDED PYROCLASTIC FACIES	181
<i>Fiamme in Welded Ignimbrites</i>	182
<i>Welded Pyroclastic Fall Deposits</i>	183
7.4 FIAMME IN SECONDARY WELDED PUMICE-RICH FACIES	184

7.5 FIAMME IN NON-WELDED, PUMICEOUS, VOLCANICLASTIC FACIES	185
<i>Submarine Syn-eruptive Gravity Current Deposits</i>	186
<i>Submarine Water-settled Pumice Deposits</i>	187
<i>Non-welded Subaerial Pyroclastic Fall Deposits</i>	189
7.6 FIAMME IN LAVAS	189
<i>Welded Autobreccia</i>	190
<i>Pumiceous Glassy vs. Dense Glassy Domains in Coherent Lava</i>	191
<i>Crystalline vs. Dense Glassy Domains in Coherent Lava</i>	192
<i>Patchy Two-phase Alteration of Coherent Lava</i>	193
7.7 DISCUSSION	195
7.8 CONCLUSIONS	196
 CHAPTER 8. GEOCHEMISTRY, GEOCHRONOLOGY AND TECTONIC SIGNIFICANCE OF THE URAL VOLCANICS AND MOUNT HOPE VOLCANICS	
8.1 INTRODUCTION	197
8.2 GEOCHRONOLOGY OF THE URAL AND MOUNT HOPE VOLCANICS	199
<i>Methods</i>	199
<i>Results</i>	203
8.3 WHOLE ROCK GEOCHEMISTRY	210
<i>Methods</i>	210
<i>Results</i>	211
<u><i>Compositions of Felsic Units</i></u>	211
<u><i>Comparison of UV and MHV Suites</i></u>	215
<u><i>Geochemical Classification and Magmatic Affinity</i></u>	217
<u><i>Mafic-intermediate Rocks</i></u>	222
8.4 TECTONIC SETTING AND PETROGENESIS	225
8.5 SUMMARY	226
 CHAPTER 9. PALAEOGEOGRAPHIC SYNTHESIS, SUMMARY AND FUTURE RESEARCH	
9.1 INTRODUCTION	228

9.2 PALAEOGEOGRAPHIC RECONSTRUCTION	228
<i>Ordovician-Devonian</i>	228
<i>Late Silurian Felsic Volcanism – the Ural Volcanics and Mount Hope Volcanics</i>	230
<i>Devonian-Carboniferous Events in the Lachlan Orogen</i>	233
9.3 ANALOGUE(S) TO THE UV AND MHV SUCCESSIONS	233
<i>Analogue A: Submarine, felsic lavas</i>	235
<i>Analogue B: Felsic volcanic successions in extensional, intra-plate continental settings</i>	236
<u><i>Okinawa Trough</i></u>	236
<u><i>Northeastern Margin of Gondwanaland</i></u>	239
<i>Analogue C: The Iberian Pyrite Belt</i>	241
<i>Discussion</i>	243
9.4 SUMMARY	244
9.5 FUTURE RESEARCH	247
REFERENCES	250
APPENDIX I	273
A. ANALYTICAL TECHNIQUES	273
I.1 XRF – University of Tasmania	273
<i>Instrument</i> <i>Philips PW1480 X-Ray Spectrometer</i>	273
<i>Sample Preparation</i>	273
<i>Corrections</i>	273
<i>Calibration</i>	274
I.2 ICPMS – University of Tasmania	274
B. STANDARDS (ALS-Chemex and UTAS)	275
APPENDIX II - GEOCHEMISTRY ANALYSES RESULTS	284
APPENDIX III - SAMPLE LIST	319

LIST OF FIGURES

CHAPTER 1

Figure 1.1 Location of the UV and MHV in central NSW and the main Lachlan Orogen	1
Figure 1.2 Landscape of the Ural Range study area	6
Figure 1.3 Landscape of the Mount Hope study area	7

CHAPTER 2

Figure 2.1 Subdivisions of the Tasman Orogenic System	11
Figure 2.2 Eastern margin of Gondwanaland in the Ordovician (~490-460 Ma)	14
Figure 2.3 The Benambran and Quidongan orogenies, Lachlan Orogen, NSW	15
Figure 2.4 Ordovician-Silurian Lachlan Orogen tectonic models	16
Figure 2.5 The Bowring-Bindi Orogeny and Cobar Deformation, Lachlan Orogen NSW	17
Figure 2.6 Siluro-Devonian Cobar Deformation (~410-390 Ma)	19
Figure 2.7 Distribution of I-, S- and A-type granitoids and A-type volcanic rocks, NSW and Victoria	20
Figure 2.8 The Tabberabberan Orogeny, Lachlan Orogen, NSW	24
Figure 2.9 Middle Devonian-Carboniferous contraction, Cobar Basin and associated basins	24
Figure 2.10 The Kanimblan Orogeny, Lachlan Orogen, NSW	25
Figure 2.11 Distribution of Early Silurian-Early Devonian volcanic successions and coeval submarine basins in southeastern Australia	29
Figure 2.12 Location of the UV and MHV within map sheet areas, NSW	33
Figure 2.13 Time-space plot of Wagga Group, Walters Range Group and Rast Group stratigraphy, Cargelligo 1:250 000 map sheet	34
Figure 2.14 Time-space plot of Mount Allen 1:100 000 map sheet stratigraphy	39
Figure 2.15 Geology of the Mount Allen, Kilparney, Lachlan Downs and Nymagee 1:100 000 map sheet areas	41

CHAPTER 3

Figure 3.1 Geological map and cross-sections of the Ural Range study area	44
Figure 3.2 Equal-area stereographic projections for the UV in the Ural Range	47
Figure 3.3 Photomicrograph of stylolitic, S_1 cleavage in 1d dacite	48
Figure 3.4 Photomicrographs of relic perthite	50
Figure 3.5 Pseudoclastic and false matrix-supported textures in coherent facies	52
Figure 3.6 Blocky, columnar jointing in 16rb rhyolite, UV	58
Figure 3.7 Felsic and mafic-intermediate coherent facies	64
Figure 3.8 Common textures in felsic coherent facies, UV	65
Figure 3.9 Monomictic breccia facies	66
Figure 3.10 Graphic log of southern section	67
Figure 3.11 Siltstone-matrix monomictic breccia facies	69
Figure 3.12 Graphic log of Browns Hill	69
Figure 3.13 Graphic log of western Mount Bowen	73
Figure 3.14 Rhyolite-dacite-siltstone breccia facies	76
Figure 3.15 Fiamme-siltstone breccia facies	78
Figure 3.16 Pumice-rich facies association	81
Figure 3.17 Graphic log of eastern "Simones Cliff" area	85
Figure 3.18 Tongues of 15rb rhyolite (r) in tan, foliated siltstone (s), north side Ural Range	88

CHAPTER 4

Figure 4.1 Schematic cross-section through the northern effusive volcanic centre and the southern explosive volcanic centre east of "Simones Cliff"	94
Figure 4.2 Geological Map of the Ural Range showing distribution of lavas and intrusions	96
Figure 4.3 Geometry and contact relationships of facies in the felsic coherent facies association of the UV	99
Figure 4.4 Facies model for the pumice-rich facies association of the UV	104
Figure 4.5 Distribution of the pumice-rich facies association, and approximate location of the inferred explosive volcanic vent	107
Figure 4.6a-d Schematic cross-sections showing the sequence of emplacement of UV facies in the Ural Range	110
Figure 4.7 Schematic facies model for a submarine, dome-top rhyolitic tuff cone, Bunga Beds and Tulumán Volcano	113

CHAPTER 5

Figure 5.1 Generalised geology and location map, Mount Hope area	117
Figure 5.2 Facies and facies associations in the Mount Hope area	118
Figure 5.3 Equal-area stereographic projections for facies in the Mount Hope area	119
Figure 5.4 Amygdale filled with green biotite and quartz. MHmv rhyolite, Mount Victor	120
Figure 5.5 Columnar jointing in MHwi rhyolite, Mount Wilson	127
Figure 5.6 Felsic Coherent Facies Association, MHV	129
Figure 5.7 Sieve-textured feldspar, MHca rhyolite, Mount Wilson	130
Figure 5.8 Unnamed Peak - graphic log of fiamme breccia facies association	131
Figure 5.9 Quartz-feldspar intergrowth textures	132
Figure 5.10 Common textures in felsic coherent facies, MHV	134
Figure 5.11 Monomictic breccia facies, Mount Victor	135
Figure 5.12 Mount Victor - graphic log and detailed map	137
Figure 5.13 Clast-bearing rhyolite facies, Mount Victor	138
Figure 5.14 Fiamme breccia facies association	143
Figure 5.15 Coan Gonn peak - graphic log and cross-section	144
Figure 5.16 Rhyolite-pumice breccia facies association	148
Figure 5.17 Mount Wilson - graphic log and detailed map	149
Figure 5.18 South Mount Wilson - graphic log	150
Figure 5.19 East Mount Allen - graphic log	152
Figure 5.20 West Mount Allen - graphic log	155
Figure 5.21 Polymictic volcanic-sedimentary conglomerate facies	156
Figure 5.22 Non-volcanic sedimentary facies association	157
Figure 5.23 Boolahbone section - graphic log	158
Figure 5.24 Nombiginni Peak, graphic log	160
Figure 5.25 Mafic-intermediate coherent facies	162
Figure 5.26 Plutonic facies	164

CHAPTER 6

Figure 6.1 Schematic diagram of the facies architecture of volcanic, non-volcanic sedimentary and plutonic facies in the Mount Hope area	169
Figure 6.2 Schematic diagram showing the distribution of syneruptive, pyroclastic facies in the southern Mount Hope area	170
Figure 6.3a-c Schematic cross sections showing the sequence of emplacement of the MHV facies in the Mount Hope area	176

CHAPTER 7

Figure 7.1	Owharoa Ignimbrite, Coromandel Volcanic Zone, North Island, New Zealand	183
Figure 7.2	Fiamme in pumice-lithic breccia in the Berserker Beds, Mt. Chalmers, QLD	185
Figure 7.3	Fiamme-siltstone breccia, UV	188
Figure 7.4	Ngongotaha Rhyolite, North Island of New Zealand	191
Figure 7.5	Pumiceous and glassy domains in coherent rhyolite, Roche Rossi Rhyolite, Italy	192
Figure 7.6	Crystalline and glassy domains in Roche Rossi Rhyolite, Italy	193
Figure 7.7	Effects of patchy, two-phase alteration on coherent rhyolite (10ra), UV	194

CHAPTER 8

Figure 8.1	Geochemistry and Geochronology sample locations, MHV	200
Figure 8.2	Geochemistry and Geochronology sample locations, UV	201
Figure 8.3	Reverse concordia diagrams of U/Pb data, UV and MHV	206
Figure 8.4	Cathode-luminescence images of zircons from UV and MHV samples	207
Figure 8.5	Harker variation diagrams of major element oxides vs. SiO_2 , UV and MHV	212
Figure 8.6	Discrimination diagrams, UV and MHV	214
Figure 8.7	SiO_2 vs. trace elements, UV and MHV	216
Figure 8.8	REE Spider plots, UV and MHV	218
Figure 8.9	Eu/Eu* vs. SiO_2 and $(\text{La}/\text{Sm})_n$ vs. SiO_2 diagrams, UV and MHV	219
Figure 8.10	A-type magma discrimination diagrams, UV and MHV	220
Figure 8.11	Additional A-type magma discrimination diagrams	222
Figure 8.12	Trace elements in UV mafic-intermediate samples	224
Figure 8.13	REE Spider plots, UV mafic-intermediate samples	224

CHAPTER 9

Figure 9.1	Schematic reconstruction of the UV during the Late Silurian-Early Devonian	231
Figure 9.2	Location of the Okinawa Trough in the East China Sea	238
Figure 9.3	Map of the northeastern margin of Gondwanaland ~70 Ma	240
Figure 9.4	Location map of the Iberian Pyrite Belt (IPB), southern Portugal and Spain	242

LIST OF TABLES

CHAPTER 2

Table 2.1 A-type Igneous Rocks of the Lachlan Orogen	23
Table 2.2 L. Silurian to E. Devonian Volcanic Successions of the Lachlan Orogen, NSW	30
Table 2.3 Summary of Stratigraphy of the Mount Hope Group	42

CHAPTER 3

Table 3.1 Facies and facies associations in the UV	54
Table 3.2 Coherent Facies in the UV	60

CHAPTER 4

Table 4.1 Estimate of unit volumes, UV lavas and intrusions	97
---	----

CHAPTER 5

Table 5.1 Facies and facies associations in the Mount Hope area	122
Table 5.2 Coherent facies in the MHV	123

CHAPTER 8

Table 8.1 LA-ICPMS U/Pb analyses results, UV and MHV samples	204
--	-----